

Mercury Concentration of Muscle Tissue and Relationship with Size of Yellowfin Tuna, *Thunnus albacares*, of the Indian Ocean

B.K.K.K. Jinadasa^{1*}, G.D.T.M. Jayasinghe¹, E.M.R.K.B. Edirisinghe² and I. Wickramasinghe³

¹Analytical Chemistry Laboratory (ACL), Institute of Post-Harvest Technology (IPHT), National Aquatic Resources Research and Development Agency (NARA), Crow Island, Colombo-15,
Sri Lanka

jinadasa76@gmail.com, thilini.madurangika123@gmail.com

² Department of chemical sciences, Faculty of Applied Sciences, Rajarata University of Sri Lanka, Mihintale,
Sri Lanka

ranjith_e@hotmail.com

³Department of Food Science and Technology, Faculty of Applied Sciences, University of Sri Jayewardenepura, Nugegoda, Sri
Lanka

indiraw2002@yahoo.com

Abstract: Mercury (Hg) is a naturally occurring metal in the earth's crust and can enter the aquatic environment through natural and anthropogenic activities. Part of Hg is converted to methyl-mercury (MeHg) and accumulates in fish through the food chain reaching its highest levels in large predatory fish such as tuna. Consumption of contaminated fish has been considered a serious public health concern. Yellow fin tuna (*Thunnus albacares*, YFT) comprises the most important component of the Indian Ocean tuna catches and it can contain significant levels of MeHg. For better understanding and monitoring purpose of Hg levels in YFT populations, total Hg (T-Hg) concentrations were analyzed in edible muscle tissue from 140 YFT collected from major fish landing sites of Sri Lanka in 2010 and 2011. The samples were analyzed using cold vapour atomic absorption spectrophotometric method, with microwave assisted digestion. In Sri Lankan waters, Hg levels in YFT ranged from <LOD (0.021) to 0.98 mg/kg (mean \pm SD = 0.30 \pm 0.18 mg/kg; median = 0.27 mg/kg) in wet weight basis. Data from the present study suggest that Sri Lankan YFT contain lower levels of Hg compared with the EU/EC recommendations (1 mg/kg). T-Hg levels of YFT were positively related with fish length and weight.

Keywords: Total mercury, Yellowfin tuna, Indian Ocean, Sri Lanka

1. Introduction

Fish and seafood products are nutrient-rich foods containing proteins, omega 3 fatty acids, vitamins, minerals etc. It's associated with various beneficial health effects and regarded as an important component of a healthy diet. Food based nutritional guidelines of many countries, including Sri Lanka; recommend the increased intake of fish and seafood. The fish consumption per capita (including dried fish and canned fish) is 15.3 kg/year, but Medical Research Institute

(MRI) of Sri Lanka recommends the increase in the per capita consumption of fish up to 21 kg/year for a healthy population [1]. However, some fish species can also contain harmful substance such as persistent organic pollutants (POPs), polychlorinated biphenyls (PCBs), dioxins, furans and other environmental contaminants such as mercury, cadmium and lead[2-5]. These contaminants are present at low levels in water systems, but bio-concentrate from one trophic level to another and reaching their highest levels in large and old predatory fish species and marine mammals [3, 6].

Mercury is a naturally occurring trace metal that is present in air, water and soil. It is an environmental pollutant that exists in several forms: elemental or metallic mercury, inorganic mercury and organic mercury. Elemental mercury has long been used in thermometers, mirrors and first aid kits. The amalgam used in dental fillings, consisting of about 50% mercury, is an example of a metallic mercury compound. Organic mercury is found mainly in fish and seafood as methylmercury species such as CH_3Hg^+ this is the most toxic forms of Hg and have received considerable attention in the world [5, 7]. Mercury contamination varies by fish species and cannot be removed through the trimming, skinning and cooking process, because MeHg binds to proteins in the muscle tissue of fish rather than to fatty deposits [8, 9]. The urine and faeces are the main excretory pathways of the elemental and inorganic Hg forms from the body with a half-life of approximately 1-2 months (WHO, 2003). Regular consumption of seafood may result in the accumulation of mercury in the body, especially when the rate of consumption exceeds the rate at which the body eliminates it. Exposure to high amounts MeHg can damage the central nervous systems (CNS), and the developing brains of young children and the foetus (WHO, 2003).

The food safety issues concerning Hg are managed by both advising on intake levels through the provisional tolerable weekly intake (PTWI) level and setting actual upper limits in foods. At present, the European Union (EU) has set an upper limit of Hg 1 mg/kg wet weight (ww) basis [10] for YFT. Public health regulations in the USA also prohibit consumption of fish with fillet Hg concentrations of more than 1 mg/kg, ww. In 1972, FAO/WHO joint expert committee on food additives (JECFA) has set a PTWI value for Hg for human sat 5 $\mu\text{g}/\text{kg}$ body weight [11].

Yellowfin tuna (YFT, *Thunnus albacares*) is one of the main target species of the commercial tuna fisheries in Sri Lanka and has a long history of scientific research. YFT often marketed as Ahi, belongs to the family scombridae and known as a highly migratory fish species. YFT is a one of the largest tuna fish species found in tropical and subtropical oceanic areas from worldwide oceans excluding Arctic Ocean [12]. YFT is preferred to leave mixed surface layer, which above the thermocline as well as contribute a large proportion of the world fishery. Sri Lanka is one of the oldest and most important YFT producing countries in the Indian Ocean [13, 14]. In 2010, it was reported that the total catch of YFT was 45,650 metric tonnes, meanwhile 1535 metric tonnes t was exported [15].

The objective of this study was the determination of the total Hg concentration of edible muscle of YFT and find out the correlation the Hg concentration with length and weight of YFT.

2. Materials and Methods:

Between July 2010 and March 2011, 140 YFT was sampled from the fish landing site; Trincomally, Beruwala, Galle and

Modara fisheries harbours of Sri Lanka (Fig. 1). These four fisheries harbours cover most important landing places specially operating high numbers of multi day fishing boats. The 500 g of edible muscle samples was taken from near the dorsal fin area of each fish, then immediately packed and iced. The total weight and total length of fish were recorded. The filleted samples were transported to the Analytical Chemical Laboratory, National Aquatic Resources Research and Development Agency (NARA) and store -20°C until further analysis




Fig. 1. Map of Sri Lanka showing the positions of sampled

All chemicals, including standard were analytical reagent (AR) quality or better and purchased from Sigma and Fluka chemicals, Switzerland. De-ionized water ($>18.2 \text{ M}\Omega/\text{cm}$) was used throughout the work. The all glassware and plastic ware were used first soak overnight in a liquid detergent solution in tap water. The glass/plastic ware then thoroughly rinse with tap water and then soaking 10% (v/v) HNO_3 overnight. Subsequent rinsing was performed using de-ionized water. Then glassware was oven dried and plastic ware was air dried prior to use.

A Varian 240 FS Atomic Absorption Spectrophotometer instrument (Varian, Springvale, Australia) equipped with and Cold Vapor Generation Accessory; VGA-77 was used for the Hg determination. For microwave assisted wet digestion, MASS 1500+ microwave system (CEM, Matthews, North Carolina, USA) was used.

For determination of Hg, the thawed samples of 1 g wet weight were submitted to microwave assisted wet digestion

using 10 mL concentrated nitric acid in the MASS XP 1500+ microwave system with a following program (1400 W, ramp time 15 min, 800 PSI, 200°C and holding time 10 min). The digests were diluted to a final volume of 50 mL with de-ionized water. Freshly prepared Hg standard solution (1 mL/L) was made by appropriate dilution and used for prepared working standard solution. A SnCl₂ solution was used as the reductant and distilled water used as an acid solution to cold vapour VGA-AAS. Blanks, samples and quality control samples were analysed following the same procedure. The Hg was assessed for compliance with the methods performance criteria guideline published by the EU, 2001/22/EC. Average recovery, precision, specificity, the detection limit and accuracy in the test were good agreement with the requirement. The trueness of Hg determination was evaluated by the analysis of the standard quality control sample, canned fish muscle T/0774 from Food Analysis Performance Assessment Scheme (Fapas), UK (n=10, certified value = 19.9 mg/kg, recovery was 98.9%) and meanwhile participated proficiency testing

program with Fapas, were used for quality control of study (canned fish, 07115/2009, Z=0. 1). The Limit of Detection (LOD = mean blank + 3s) and Limit of quantification (LOQ =3 x LOD) for Hg was estimated from the blank measurements according to EURACHEM, and the value was 0.007 mg/kg and 0.021 mg/kg respectively.

The statistical analysis was performed using Microsoft Excel, 2011, SPSS (Statistical Package for Social Sciences) 16 and Curve Expert Professional software.

3. Results and Discussion:


This study provides information on T-Hg levels in the muscle tissue of YFT collected from the landing sites of Sri Lanka, and also examines the relationship between T-Hg concentration with length and weight of fish. A total of 140 YFT were analysed and results are given in table 1.

Table 1: Length, weight measurement and mercury content of 140 samples of YFT

	Weight (kg)	Length (cm)	T-Hg (mg/kg), w/w
Avg. ± SD	45.28 ± 13.96	123.4 ± 23.8	0.30 ± 0.18
Range	18.00 – 83.50	64.0 - 173.0	<LOD (0.021) – 0.98

Figure 2 shows the correlations between YFT weight, length and T-Hg content. There is a positive correlation with weight of fish ($R^2=0.24$, $r=0.53$) for larger fish to have a higher Hg content. The model fit is linear one in which $Y=a + bX$ where $a = 0.9688$, $b= 3.742$ and standard error was 0.09. This slight correlation is also shown in Figure 2 with the correlation between YFT length and T-Hg content ($R^2=0.15$, $r=0.41$) which is a linear model fit. The model values were; $a = -5.976$,

$b = 2.463$ and standard error = 0.152. The linear relationship between age or size of many carnivore species and Hg concentration is well documented and this study also confirms that. By the calculation of linear regression equation, it was found that, the Hg levels in YFT weighing more than 249 kg or above length of 408 cm, exceed the 1.0 mg/kg level. Nevertheless, YFT of such length and weight was not recorded during the present study.






Fig. 2: Correlation between weight, length and mercury concentration of YFT in the present study

In comparison with published data, the Hg levels detected in YFT during this study were quite similar to YFT from oceans around the world as shown in table 2.

Table 2: T-Hg (mean and range) of YFT recorded in other studies

T-Hg (mg/kg)	T-Hg range (mg/kg)	Area	Reference
	0.026-0.234	Andaman Sea	[16]
0.25	0.068-0.650	Florida Atlantic coast	[17]
0.092	0.061-0.124	Bay of Bengal	[18]
0.21	–	Pacific ocean	[19]
0.13	–	Mozambique Channel	[20]
0.21	–	Reunion Island	[20]
0.54	0.24-1.32	Hawaii	[21]
0.42	0.07-1.20	Gulf of Guinea	[21]
0.05	0.01-0.08	West coast, Sri Lanka	[22]
0.30	<0.021-0.98	Indian Ocean, around Sri Lanka	This study

Research and monitoring of the bio-geochemical cycle of Hg in the marine environment is critical to expanding the current understanding of the Hg sources that contaminated the fish. Limited information exists regarding T-Hg in YFT from the Indian Ocean around Sri Lanka. In an earlier study by Senadheera, 2005, 40 muscle samples of YFT collected from the west coast in Sri Lanka were analysed. It is not clear if the total length of the fish was studied, but the mean weight was 48 kg. T-Hg of these fish ranged from 0.01-0.08 mg/kg in wet weight basis. The mean T-Hg value of that

study (0.05 mg/kg) and the range of Hg concentration were lower than those observed in this study, but sampling procedure and analytical method used in the two studies may not be directly comparable. For that study, the samples were collected from a local fisherman on the west coast and from fish export factory, as well as for Hg analyses was performed by the colorimetric dithizone method described in Association of Analytical Communities (Association of Official Analytical Chemists, AOAC) 952.14.

Several factors are likely to be responsible for the observed differences in T-Hg levels in YFT. For marine fishes, dietary exposure, size and age of fish and trophic levels are “important sources of T-Hg” (Joanna and Michael, 2006 and Wang, 2002). Other factors are temperature, salinity, pollution sources, sexual state, feeding mechanisms and physiological state of the animals [23]. In general, larger fish contain higher T-Hg concentrations than smaller fish within the species [24]. Fish are in a higher position in the aquatic food web and therefore they accumulate more biomagnifying contaminants such as Hg [25]. YFT principally consumes a variety of crustaceans and squids (www.fishbase.org). YFT is classified as a highly migratory fish species by FAO 1997, and the difference of T-Hg may be prey available during the migration. Information regarding the age and growth of YFT in Indian Ocean surrounding Sri Lanka is limited, and specific age of species in this study has not been determined. The preliminary estimate of the age of YFT from this study based on length and weight data from other regions, indicate that of fish samples are between 1.5 and 4+ years old. The mean growth rate of the YFT in Indian Ocean was approximately 1-2 mm/day [26].

Fish and seafood are highly regarded as rich in nutrition, especially protein, omega 3 fatty acids, vitamins and minerals, thus challenges still remain with regard to balancing risk benefit messages regarding fish consumption and Hg. Predatory marine fish species, especially in upper trophic levels, such as YFT can accumulate relatively high levels of Hg. The nature of bioaccumulation of methyl Hg based on several factors including multiple anthropogenic and natural sources. Methyl Hg exposure through fish consumption will continue to remain as a public health concern. The consumption of fish and fishery products is the main pathway for human exposure to Hg [2]. Thus,

References

1. NARA, *Sri Lanka fisheries yearbook*. 2009, NARA: Colombo, Sri Lanka.
2. Gholam, R.J.K., et al., *Mercury contamination in fish and public health aspects: A review*. Pakistan journal of nutrition, 2005. **4**(5): p. 276-281.
3. Roser, M.C., et al., *Intake of chemical contaminants through fish and seafood consumption by children of Catalonia, Spain: Health risks*. Food and Chemical Toxicology, 2007. **45**(10): p. 1968-1974.
4. Silvia, T.E., V. Dinoraz, and M. Rosa, *Mercury and methyl mercury bioaccessibility in swordfish*. Journal of food additives and contaminants, 2010. **27**(3): p. 327-337.
5. Jyrki, K.V., et al., *Mercury as a risk factor for cardiovascular diseases*. Journal of nutritional biochemistry, 2006. **18**(2): p. 75-85.
6. Ross, P.S., et al., *High PCB concentrations in free ranging Pacific Killer Whales, Orcinus orca: Effects of age, sex and dietary preference*. Marine Pollution Bulletin, 2000. **40**(6): p. 504-515.
7. Milena, H., et al., *Total mercury, methylmercury and selenium in mercury polluted area in the province Guizhou, China*. The science of the total environment, 2003. **304**: p. 231-256.
8. Domingo, J.L., et al., *Benefits and risk of fish consumption, II RIBEPEIX, a computer program to optimize the balance between the intake of omega-3 fatty acid and chemical contaminants*. Toxicology, 2007. **230**: p. 227-233.
9. FAO, *Health risks associated with fish consumption focus on methyl-mercury, dioxins and dioxin like PCB'S*. 2010, FAO: Rome, Italy.
10. EU/EC-1881, *Commission Regulation (EC), No 1881/06 of setting maximum levels for certain contaminants in foodstuffs*. Official Journal of European Union, 2006. **L364**: p. 5-24.

4. Conclusions

It can be concluded that the T-Hg content of the Indian Ocean around Sri Lanka, as reported in this study are well bellowed EU/EC regulation tolerance limits of 1.0 mg/kg. It is recommended that population groups at risk in Sri Lanka should limit consumption of YFT and consumption of large YFT should be avoided.

Acknowledgement

The authors thank the National Aquatic Resources Research & development Agency (NARA), project No. 13.2, 2009 to provide funds and the authorization to publish this work and the Fish exporters and Mr. M.H.S.K. Abeyrathne for help in sampling and analysis.

11. FAO/WHO, *Evaluataion of the certain food additives and the contaminents mercury, lead and cadmium*. 1972, World health organization, technical report series No. 505.
12. FAO. *Fisheries Data bases and Statistics*. 2009 [cited 2010 11/22].
13. Disanayake, D.C.T., E.K.V. Samaraweera, and C. Amarasiri, *Fishery and feeding habits of Yellowfin tuna (Thunnus albacores) targeted by coastal tuna long lining in the north western and north eastern coast of Sri Lanka*. Sri Lanka journal of aquatic science, 2008. **13**: p. 1-21.
14. Rajapaksha, J.K., T. Nishida, and L. Samarakoon. *Environmental preferences of yellow fin tuna (Thunnus albacores) in the northeast Indian Ocean: an application of remote sensing data to long-line catches*. in *IOTC Fourteenth Working Party on Tropical Tunas*. 2010. Mauritius: IOTC.
15. MOFAR, *Ministry of fisheries and aquatic reources, fisheries statistics, Sri Lanka, Accessed date 2013-01-10*. 2011.
16. Menasveta, P. and R. Siriyong, *Mercury content of several predacious fish in the Andaman Sea*. Marine pollution bulletin, 1976. **8**: p. 200-204.
17. Adams, D.H., *Total mercury levels in tunas from offshore waters of the Florida Atlantic coast*. Marine pollution bulletin, 2004. **49**: p. 659-667.
18. Penjai, S., et al., *An assessment of mercury concentration in fish tissues caught from three compartments of Bay of Bengal, in The ecosystem based fishery management in the Bay of Bengal*. 2008, BOBLME. p. 221-232.
19. Kraepiel, A.M.L., et al., *Sources and variations of mercury in tuna*. Environmental science and technology, 2003. **37**: p. 5551-5558.
20. Jessica, K., et al., *Mercury content in commercial pelagic fish and its risk assessment in the Western Indian Ocean*. Journal of science of the total environment, 2006. **366**(2-3): p. 688-700.
21. Peterson, C.L., W.L. Klawe, and G.D. Shar, *Mercury in tunas: A review*. Fisheries bulletin, 1973. **71**(3): p. 603-613.
22. Senadheera, S.P.S.D., *Fish for human consumption: is there a risk of contamination with mercury?* Journal of the national aquatic resources & research development agency of Sri Lanka, 2005. **37**: p. 61-68.
23. El-Moselhy, K.-M., *Bioaccumulation of mercury in some organisms from Lake Timsah and Bitter Lake (Suez Canal, Egypt)*. Egyptian journal of aquatic research, 2006. **32**(1): p. 124-134.
24. Joanna, B. and G. Michael, *Mercury in fish available in supermarkets in Illinois: Are there regional differences*. Science of the total environment, 2006. **367**: p. 1010-1016.
25. Campbell, K.R., *Concentrations of heavy metals associated with urban runoff in fish living in storm water treatment ponds*. Archives of environmental contamination and toxicology, 1994. **27**: p. 352-356.
26. Bernard, S. and C. Francois, *Age and growth of big eye tuna (Thunnus obessus) in Western Indian Ocean*. Cybium, 1996. **28**(2): p. 163-170.
27. MOFAR, *Gazette No. 1528/7, The Fish Products (Export) Regulations, Act No 2 of 1996*. 2007, Ministry of Fisheries and Aquatic Resources: Sri Lanka.